

# Solutions

Name:

## Math 107H Exam 1

Show all work. How you get your answer is just as important, if not more important, than the answer itself.

Find each of the following integrals.

Note that " $\int_3^x f(t) dt + C$ " is not a sufficient computation of an antiderivative!

Some formulas of potential use can be found at the bottom of the last page of the exam.

1. (10 pts.)  $\int (x+2)^{3/2} dx$

$$\begin{aligned} u &= x+2 \quad du = dx \\ \int (x+2)^{3/2} dx &= \int u^{3/2} du \Big|_{u=x+2} = \frac{2}{5} u^{\frac{5}{2}} + C \Big|_{u=x+2} \\ &= \frac{2}{5} (x+2)^{\frac{5}{2}} + C \end{aligned}$$

3 is odd!

2. (15 pts.)  $\int_0^{\pi/2} \sin^3 x \, dx$

$$= \int_0^{\frac{\pi}{2}} \sin^2 x (\sin x \, dx)$$

$$= \int_0^{\frac{\pi}{2}} (1 - \cos^2 x) (\sin x \, dx)$$

$$u = \cos x \quad du = -\sin x \, dx$$

$$x=0 \rightsquigarrow u=1$$

$$x=\frac{\pi}{2} \rightsquigarrow u=0$$

$$= \int_1^0 (1 - u^2) (-du) = \int_1^0 u^2 - 1 \, du$$

$$= \left. \frac{u^3}{3} - u \right|_1^0 = \boxed{(0 - 0) - \left( \frac{1}{3} - 1 \right)}$$

$$= 0 - \left( -\frac{2}{3} \right) = \frac{2}{3}$$

$$\begin{aligned}
 3. \text{ (10 pts.)} \int \frac{x^2 + x - 3}{x^{1/2}} dx &= \int \frac{x^2}{x^{1/2}} + \frac{x}{x^{1/2}} - \frac{3}{x^{1/2}} dx \\
 &= \int x^{3/2} + x^{1/2} - 3x^{-1/2} dx \\
 &= \left[ \frac{2}{5}x^{5/2} + \frac{2}{3}x^{3/2} - 3(2x^{1/2}) + C \right]
 \end{aligned}$$

$$4. (15 \text{ pts.}) \int_0^1 e^{\sqrt{x}} dx$$

$u = \sqrt{x} \quad du = \frac{1}{2}x^{-\frac{1}{2}} dx = \frac{1}{2\sqrt{x}} dx$

$$u=0 \rightarrow x=0$$

$$u=1 \rightarrow x=1$$

$$= \int_0^1 (2\sqrt{x}) e^{\sqrt{x}} \left( \frac{dx}{2\sqrt{x}} \right) = \int_0^1 2ue^u du$$

$$\begin{aligned} v &= u \quad dw = e^u du \\ dv &= du \quad w = e^u \end{aligned} \quad = 2 \int_0^1 ue^u du$$

$$= 2 \left( ue^u \Big|_0^1 - \int_0^1 e^u du \right)$$

$$= 2 \left( ue^u \Big|_0^1 - e^u \Big|_0^1 \right)$$

$$\boxed{2((1 \cdot e - 0 \cdot 1) - (e - 1))}$$

$$= 2(1 - e + 1) = 2(1) = \underline{\underline{2}}$$

$u = e^{\sqrt{x}}$  also works! Gets you to  $2 \int u du \Big|_{u=e^{\sqrt{x}}}$ ...

$$5. \text{ (15 pts.) } \int \frac{dx}{(x+1)^2(x+4)} = (\star)$$

$$\begin{aligned} \frac{1}{(x+1)^3(x+4)} &= \frac{A}{x+1} + \frac{B}{(x+1)^2} + \frac{C}{(x+4)} \\ &= \frac{A(x+1)(x+4) + B(x+4) + C(x+1)^2}{(x+1)^2(x+4)} \end{aligned}$$

so need:  $1 = A(x+1)(x+4) + B(x+4) + C(x+1)^2$

Set  $x=-1$ :  $1 = A(-1)(3) + B(3) + C(-1)^2 = 3B$   
 $\Rightarrow B = \frac{1}{3}$

Set  $x=-4$ :  $1 = A(-3)(0) + B(0) + C(-3)^2 = 9C$   
 $\Rightarrow C = \frac{1}{9}$

Set  $x=0$ :  $1 = A(1)(4) + \frac{1}{3}(4) + \frac{1}{9}(1)^2 = 4A + \frac{4}{3} + \frac{1}{9}$   
 $\Rightarrow 4A = 1 - \frac{4}{3} - \frac{1}{9} = \frac{9-12-1}{9} = -\frac{4}{9} \Rightarrow A = -\frac{1}{9}$

so  $(\star) = \int -\frac{1}{9} \frac{1}{x+1} + \frac{1}{3} \frac{1}{(x+1)^2} + \frac{1}{9} \frac{1}{x+4} dx$

$(u=x+1) \quad (u=x+1) \quad (u=x+4)$   
 $(du=dx) \quad (du=dx) \quad (du=dx)$

$$\boxed{\int -\frac{1}{9} \ln|x+1| + \frac{1}{3} \left( \frac{-1}{(x+1)} \right) + \frac{1}{9} \ln|x+4| + C}$$

6. (15 pts.)  $\int e^{-x} \sin(3x) dx$

(1)

$$u = \sin(3x) \quad dv = e^{-x} dx$$

$$du = 3 \cos(3x) dx \quad v = -e^{-x}$$

$$= -e^{-x} \sin(3x) - \int -3e^{-x} \cos(3x) dx$$

$$= -e^{-x} \sin(3x) + 3 \int e^{-x} \cos(3x) dx$$

$$u = \cos(3x) \quad dv = e^{-x} dx$$

$$du = -3 \sin(3x) \quad v = -e^{-x}$$

$$= -e^{-x} \sin(3x) + 3 \left( -e^{-x} \cos(3x) - \int -(-3)e^{-x} \sin(3x) dx \right)$$

$$= -e^{-x} \sin(3x) - 3e^{-x} \cos(3x) - 9 \int e^{-x} \sin(3x) dx$$

S  $10 \int e^{-x} \sin(3x) dx = -e^{-x} \sin(3x) - 3e^{-x} \cos(3x)$

so:  $\int e^{-x} \sin(3x) dx = -\frac{1}{10} e^{-x} \sin(3x) - \frac{3}{10} e^{-x} \cos(3x) + C$

$$7. (20 \text{ pts.}) \int (x^2 + 1)^{3/2} dx = \int (\sqrt{x^2 + 1})^3 dx$$

Think:  $\tan^2 u + 1 = \sec^2 u$

$$x = \tan u \quad dx = \sec^2 u du$$

~~$$dx^2 \quad x^2 + 1 = \tan^2 u + 1 = \sec^2 u$$~~

$$\sqrt{x^2 + 1} = \sec u$$

$$= \int (\sec u)^3 (\sec^2 u du) \Big|_{x=\tan u}$$

$$= \int \sec^5 u du \Big|_{x=\tan u} \stackrel{(n=5)}{=} \frac{1}{4} \sec^3 u \tan u \Big|_{x=\tan u} + \frac{3}{4} \int \sec^3 u du \Big|_{x=\tan u}$$

$$\stackrel{(n=3)}{=} \frac{1}{4} \sec^3 u \tan u + \frac{3}{4} \left( \frac{1}{2} \sec u \tan u + \frac{1}{2} \int \sec u du \right) \Big|_{x=\tan u}$$

$$= \frac{1}{4} \sec^3 u \tan u + \frac{3}{8} \sec u \tan u + \frac{3}{8} \ln |\sec u + \tan u| + C \Big|_{x=\tan u}$$

$$= \frac{1}{4} (\sqrt{x^2 + 1})^3 x + \frac{3}{8} (\sqrt{x^2 + 1}) x + \frac{3}{8} \ln |\sqrt{x^2 + 1} + x| + C$$

$\sqrt{x^2 + 1}$   $x \rightarrow \tan u = x$   
 $\sec u = \sqrt{x^2 + 1}$

$$\int \sec^n x dx = \frac{1}{n-1} \sec^{n-2} x \tan x + \frac{n-2}{n-1} \int \sec^{n-2} x dx$$

$$c^2 \int \frac{dy}{(y^2 + c^2)^k} = \frac{1}{(2k-2)} \cdot \frac{y}{(y^2 + c^2)^{k-1}} + \frac{(2k-3)}{(2k-2)} \int \frac{dy}{(y^2 + c^2)^{k-1}}$$

1. Find the following integrals (10 pts. each):

$$(a): \int_1^4 x^2 \ln x \, dx \quad \text{by parts!} \quad u = \ln x \quad dv = x^2 dx \\ du = \frac{1}{x} dx \quad v = \frac{1}{3} x^3$$

$$= \frac{1}{3} x^3 \ln x \Big|_1^4 - \int_1^4 \frac{1}{3} x^3 dx = \frac{1}{3} x^3 \ln x \Big|_1^4 - \frac{1}{9} x^3 \Big|_1^4 \\ = \left( \frac{1}{3} 4^3 \ln 4 - \frac{1}{3} 1^3 \ln 1 \right) - \frac{1}{9} (4^3 - 1^3)$$

[can also be done by  $u = x^2 \ln x, dv = dx$ !]

$$(b): \int \sin^2 x \cos^3 x \, dx = \int \sin^2 x \cos^2 x \cos x \, dx \\ = \int \sin^2 x (1 - \sin^2 x) \cos x \, dx \quad u = \sin x \quad du = \cos x \, dx \\ = \int u^2 (1 - u^2) du \Big|_{u=\sin x} = \int u^2 - u^4 du \Big|_{u=\sin x} \\ = \frac{1}{3} u^3 - \frac{1}{5} u^5 + C \Big|_{u=\sin x} = \frac{1}{3} (\sin x)^3 - \frac{1}{5} (\sin x)^5 + C$$

2. When you apply the appropriate trigonometric substitutions, what do the following integrals become?

$$(a): \int \frac{\sqrt{4-x^2}}{x^2} dx$$

$$x = 2\sin u \quad dx = 2\cos u du$$

$$\sqrt{4-x^2} = \sqrt{4-4\sin^2 u} = 2\cos u$$

$$= \left| \int \frac{(2\cos u)(2\cos u du)}{(2\sin u)^2} \right| = \left| \int \frac{\cos^2 u}{\sin^2 u} du \right| \Bigg|_{x=2\sin u}$$

$$(b): \int \frac{x^2}{\sqrt{4x^2+9}} dx$$

$$\text{want } 4x^2+9 = 9\tan^2 u + 9 = 9\sec^2 u$$

$$2x = 3\tan u \quad x = \frac{3}{2}\tan u$$

$$dx = \frac{3}{2}\sec^2 u du$$

$$\sqrt{4x^2+9} = \sqrt{9\sec^2 u} = 3\sec u$$

$$= \left| \int \frac{\left(\frac{3}{2}\tan u\right)^2 \left(\frac{3}{2}\sec^2 u du\right)}{3\sec u} \right| = \left| \int \frac{\frac{9}{4}\tan^2 u \sec u du}{3\sec u} \right| \Bigg|_{x=\frac{3}{2}\tan u}$$

6. (15 pts.) Recall that if a function  $f$  has second derivative satisfying  $|f''(x)| \leq M$  for every  $x$  in the interval  $[a, b]$ , then the error  $E_n$  in approximating the integral  $\int_a^b f(x) dx$  using the trapezoidal rule using  $n$  equal subintervals is at most

$$M \frac{(b-a)^3}{12n^2}$$

Based on this, how many subintervals should we divide the interval  $[2, 5]$  into in order to be sure to approximate the integral  $\int_2^5 x \ln x dx$  with an error of less than  $\frac{1}{100}$ ?

$$\begin{aligned} f(x) &= x \ln x \\ f'(x) &= 1 \cdot \ln x + x \left(\frac{1}{x}\right) = \ln x + 1 \\ f''(x) &= \frac{1}{x} + 0 = \frac{1}{x} \quad \text{← decreasing!} \quad (f'''(x) = -\frac{1}{x^2} < 0) \\ \text{so max is at } x=2 &\quad M = \frac{1}{2} \end{aligned}$$

$$\text{Want error} < \frac{1}{100}, \text{ error} \leq \frac{1}{2} \frac{(5-2)^3}{12n^2}$$

$$\Leftrightarrow \text{want } \frac{1}{2} \frac{3^3}{12n^2} < \frac{1}{100} \Rightarrow n^2 > 100 \frac{1}{2} \frac{3^3}{12} \\ = 50 \frac{27}{12} = 50 \frac{9}{4}$$

$$\text{so need } n > \sqrt{112.5} \approx 10.6 \quad = \frac{450}{9} = 112.5$$

$$\text{so } n=11 \text{ will work.}$$