


Old final solutions

H-1: $\int \sec^3 x \tan^3 x dx = \int \sec^2 x \tan^2 x (\sec x \tan x dx)$
 $= \int \sec^2 x (\sec^2 x - 1) (\sec x \tan x dx)$ $[u = \sec x \quad du = \sec x \tan x dx]$
 $= \int u^2(u^2 - 1) du \Big|_{u=\sec x} = \int u^4 - u^2 du \Big|_{u=\sec x} = \frac{u^5}{5} - \frac{u^3}{3} + C \Big|_{u=\sec x}$
 $= \boxed{\frac{1}{5} \sec^5 x - \frac{1}{3} \sec^3 x + C}$

H-2: $\int \frac{x^2 dx}{\sqrt{3-x^2}}$ $x = \sqrt{3} \sin u, dx = \sqrt{3} \cos u du, 3-x^2 = 3 \cos^2 u$
 $= \int \frac{(\sqrt{3} \sin u)^2 (\sqrt{3} \cos u du)}{\sqrt{3 \cos^2 u}} = \int 3 \sin^2 u du \Big|_{x=\sqrt{3} \sin u}$
 $= 3 \int \frac{1}{2} (1 - \cos 2u) du \Big|_{x=\sqrt{3} \sin u} = \frac{3}{2} (u - \frac{1}{2} \sin 2u + C) \Big|_{x=\sqrt{3} \sin u}$
 $= \frac{3}{2} (u - \sin u \cos u) + C \Big|_{x=\sqrt{3} \sin u}$

 $= \boxed{\frac{3}{2} (\text{Arcsin}(\frac{x}{\sqrt{3}})) - \frac{x}{\sqrt{3}} \frac{\sqrt{3-x^2}}{\sqrt{3}} + C}$

H-3: $\int x^2 e^{3x} dx$ $u = x^2, dv = e^{3x} dx \quad du = 2x dx, v = \frac{1}{3} e^{3x}$
 $= \frac{1}{3} x^2 e^{3x} - \frac{2}{3} \int x e^{3x} dx$ $u = x \quad dv = e^{3x} \quad du = dx \quad v = \frac{1}{3} e^{3x}$
 $= \frac{1}{3} x^2 e^{3x} - \frac{2}{3} (\frac{1}{3} x e^{3x} - \frac{1}{3} \int e^{3x} dx) = \boxed{\frac{1}{3} x^2 e^{3x} - \frac{2}{9} x e^{3x} + \frac{2}{27} e^{3x} + C}$

H-4: $\int \frac{2x+3}{x^3+x^2-2} dx = \int \frac{2x+3}{(x-1)(x^2+2x+2)} dx = \int \frac{A}{x-1} + \frac{Bx+C}{x^2+2x+2} dx$
 $2x+3 = A(x^2+2x+2) + (x-1)(Bx+C) \quad x=1, 5 = A(5) \rightarrow A=1$
 $x=0, 3 = (1)(2) + (-1)(C), C = 2-3 = -1, x=-1, 1 = (1)(1) + (-2)(-B-1)$
 $-B-1 = 0, B = -1$

$$\int \frac{2x+3}{x^3+x^2-2} dx = \int \frac{1}{x-1} - \frac{x+1}{x^2+2x+2} dx = \ln|x-1| - \int \frac{x+1}{(x+1)^2+1} dx$$

$$= \ln|x-1| - \int \frac{u}{u^2+1} du \Big|_{u=x+1} = \ln|x-1| - \int \frac{\frac{1}{2} dv}{v} \Big|_{v=u^2+1} \Big|_{u=x+1}$$

$$= \ln|x-1| - \frac{1}{2} \ln|v| + C \Big|_{v=u^2+1} \Big|_{u=x+1} = \ln|x-1| - \frac{1}{2} \ln|(x+1)^2+1| + C \Big|_{u=x+1}$$

$$= \ln|x-1| - \ln|(x+1)^2+1| + C$$

2. $f(x) = g(x) : 2x-1 = x^4+x-1, x^4-x = 0 = (x^3-1)x$

$\Rightarrow x=0$ or $x^3-1=0 \rightarrow x^3=1 \rightarrow x=1$


on $[0,1]$, $2x-1 \geq x^4+x-1$ (check $x=\frac{1}{2} : 0 = 1-1 > \frac{1}{16} + \frac{1}{2} - 1$)

Area = $\int_0^1 (2x-1) - (x^4+x-1) dx = \int_0^1 -x^4+x dx$

$$= \frac{x^2}{2} - \frac{x^5}{5} \Big|_0^1 = \left(\frac{1}{2} - \frac{1}{5}\right) - (0-0) = \frac{1}{2} - \frac{1}{5} = \frac{5}{10} - \frac{2}{10} = \frac{3}{10}$$

$x^3+7x-22=0 : (x-2)(x^2+2x+11)=0 \quad \underline{x=2}$

3.



By shells: (width) (height)

Volume = $\int_0^2 2\pi(x-2)(x^3+7x-22) dx$


$= \int_0^2 2\pi(x+2)(x^3+7x-22) dx = 2\pi \int_0^2 x^4+2x^3+7x^2+14x-22x-44 dx$

$= 2\pi \int_0^2 x^4+2x^3+7x^2-8x-44 dx = 2\pi \left(\frac{x^5}{5} + \frac{x^4}{2} + \frac{7x^3}{3} - 4x^2 - 44x \right) \Big|_0^2$

$= 2\pi \left(\left(\frac{2^5}{5} + \frac{2^4}{2} + \frac{7 \cdot 8}{3} - 4 \cdot 4 - 44 \cdot 2 \right) - (0) \right)$

IGNORE...

4.



$$\text{Area} = A(x) = \pi x^2 = \pi(144 - y^2)$$

$$\text{work} = \int_{-12}^0 \text{dist}(-y) \cdot \underbrace{\pi(144 - y^2)}_{\text{mass}} dy$$

$$= 300\pi \int_{-12}^0 (-144y + y^3) dy = 300\pi \left(-72y^2 + \frac{y^4}{4} \right) \Big|_{-12}^0$$

$$= 300\pi \left((0+0) - \left(-72(-12)^2 + \frac{(-12)^4}{4} \right) \right)$$

$$= 300\pi \left(72 \cdot 12^2 - \frac{12^2}{4} \cdot 12^2 \right) = 300\pi \cdot 12^2 (72 - 36)$$

$$= \boxed{300\pi \cdot 12^2 \cdot 36}$$

5 (a) $\lim_{x \rightarrow \infty} \frac{x^2 - 3x^3 + 9}{4x^2 - 6x + 1} = \lim_{x \rightarrow \infty} \frac{1 - 3x + \frac{9}{x^2}}{4 - \frac{6}{x} + \frac{1}{x^2}} \approx \frac{\text{large neg}}{4} = \boxed{-\infty}$

(b) $\lim_{x \rightarrow \infty} \frac{(x^2+1)^x}{(x+1)^{2x}} = L$ $\ln L = \lim_{x \rightarrow \infty} x \ln(x^2+1) - 2x \ln(x+1)$

$$= \lim_{x \rightarrow \infty} x (\ln(x^2+1) - 2 \ln(x+1)) = \lim_{x \rightarrow \infty} x \ln \left(\frac{x^2+1}{(x+1)^2} \right) = \lim_{x \rightarrow \infty} \frac{\ln \left(\frac{1+(\frac{1}{x})^2}{(1+\frac{1}{x})^2} \right)}{\frac{1}{x}}$$

$$= \lim_{h \rightarrow 0^+} \frac{1}{h} \ln \left(\frac{1+h^2}{(1+h)^2} \right) = f'(0), \quad f(x) = \ln \left(\frac{1+x^2}{(1+x)^2} \right)$$

But: $f'(x) = \left(\frac{1}{\frac{1+x^2}{(1+x)^2}} \right) \left(\frac{(1+x)^2(2x) - (1+x^2)(2(1+x))}{(1+x)^2} \right)$; at $x=0$,

$$f'(0) = \frac{1}{\left(\frac{1}{1^2}\right)} \left(\frac{(1)(0) - (1)(2)}{1^2} \right) = -2 \quad \therefore \ln L = -2, \quad \boxed{L = e^{-2}}$$

6-1: $\sum_{n=1}^{\infty} \frac{(n+1)^{1/2}}{n^2} = \sum a_n$ $b_n = \frac{n^{1/2}}{n^2} = \frac{1}{n^{3/2}}$ then

$\frac{a_n}{b_n} = \frac{(n+1)^{1/2}}{n^{3/2}} = \left(1 + \frac{1}{n}\right)^{1/2} \rightarrow 1 \neq 0$ as $n \rightarrow \infty$, so since

$\sum b_n$ converges (p-series, $p=3/2 > 1$), $\sum a_n$ conv by lin. compar.

6-2: $\sum \frac{n!}{(n^2+n-3)^{3/2}} = \sum a_n$ $b_n = \frac{n!}{(n^2)^{3/2}} = \frac{n!}{n^3}$ then

$\frac{a_n}{b_n} = \left(\frac{n^2}{n^2+n-3}\right)^{3/2} \rightarrow 1 \neq 0$ as $n \rightarrow \infty$. But $\sum \frac{n!}{n^3} \geq \sum \frac{n(n-1)(n-2)!}{n^3}$

and $\frac{n(n-1)(n-2)}{n^3} = (1-\frac{1}{n})(1-\frac{2}{n}) \rightarrow 1 \neq 0$ as $n \rightarrow \infty$ so $\sum b_n$ diverges by n^{th} term test, so $\sum a_n$ diverges by lin compar.

6-3: $\sum \left(\frac{n+3}{3n-5}\right)^n = \sum a_n$ $a_n^{1/n} = \frac{n+3}{3n-5} = \frac{1+3/n}{3-5/n} \rightarrow \frac{1}{3} < 1$
 as $n \rightarrow \infty$ so $\sum a_n$ conv by the root test.

6-4: $\sum_{n=1}^{\infty} \frac{\ln n}{n^{5/3}} = \sum a_n$. But $\frac{\ln n}{n^{\alpha}} \rightarrow 0$ as $n \rightarrow \infty$ for any $\alpha > 0$, so compare to $b_n = \frac{n^{1/3}}{n^{5/3}} = \frac{1}{n^{4/3}}$.

$\frac{a_n}{b_n} = \frac{\ln n}{n^{1/3}} \rightarrow 0$ as $n \rightarrow \infty$ and since $\sum b_n$ conv (p-series, $p=4/3 > 1$) $\sum a_n$ conv by limit comparison.

7: $f(x) = (x^2-5)^{5/2}$ centered at $c=3$. $f(3) = (9-5)^{5/2} = 4^{5/2} = 2^5 = 32$.
 $f'(x) = \frac{5}{2}(x^2-5)^{3/2}(2x)$ $f'(3) = \frac{5}{2}(4)^{3/2}(6) = 5 \cdot 2^3 \cdot 3 = 120$

$f''(x) = \frac{5}{2} \left(\frac{3}{2}(x^2-5)^{1/2}(2x)(2x) + 2(x^2-5)^{3/2} \right)$
 $f''(3) = \frac{5}{2} \left(\frac{3}{2} 4^{1/2}(6)(6) + 2(4)^{3/2} \right) = \frac{5}{2} \left(108 + 16 \right) = 310$

$$f'''(x) = \frac{5}{2} \left(\frac{3}{2} (x^2-5)^{-1/2} (2x)(4x^2) + 2(8x)(x^2-5)^{1/2} \right) + 2 \left(\frac{3}{2} \right) (x^2-5)^{1/2} (2x)$$

$$f'''(3) = \frac{5}{2} \left(\frac{3}{2} \left(\frac{1}{2} (4)^{-1/2} (6)(4 \cdot 6^2) + (24)(4)^{1/2} \right) + 3(4)^{1/2}(6) \right)$$

$$= \frac{5}{2} \left(\frac{3}{2} (216 + 48) + 36 \right) = \frac{5}{2} \left(\frac{3 \cdot 132 + 36}{396} \right) = \frac{5}{2} \left(\frac{432}{216} \right) = 1080$$

$$\delta \quad P_3(x) = f(3) + f'(3)(x-3) + \frac{f''(3)}{2}(x-3)^2 + \frac{f'''(3)}{6}(x-3)^3$$

$$= 32 + 120(x-3) + 155(x-3)^2 + 180(x-3)^3$$

P: $x(t) = t^4, y(t) = t^6 \quad x'(t) = 4t^3 \quad y'(t) = 6t^5$

$$\text{Length} = \int_0^2 \sqrt{(4t^3)^2 + (6t^5)^2} dt = \int_0^2 \sqrt{16t^6 + 32t^{10}} dt$$

$$= \int_0^2 t^3 (16 + 32t^4)^{1/2} dt$$

$$u = 16 + 32t^4 \quad du = 128t^3 dt$$

$$t^3 dt = \frac{1}{128} du$$

$$t=0 \rightarrow u=16$$

$$t=2 \rightarrow u = 16 + 32 \cdot 16 = 33 \cdot 16 = 480 + 16 = 528$$

$$= \int_{16}^{528} u^{1/2} \frac{1}{128} du$$

$$= \frac{1}{128} \frac{2}{3} u^{3/2} \Big|_{16}^{528}$$

$$= \frac{1}{3 \cdot 64} \left((528)^{3/2} - 16^{3/2} \right)$$

5(b), REDUX: $\frac{(x^2+1)^x}{(x+1)^{2x}} = \left(\frac{x^2+1}{(x+1)^2} \right)^x = \left(\frac{x^2+1}{x^2(x+1)} \right)^x = \left(1 - \frac{2x}{(x+1)^2} \right)^x$

$$= \left(1 - \frac{2}{\frac{(x+1)^2}{x}} \right)^x = \left(1 - \frac{2}{\frac{(x+1)^2}{x}} \right)^{\frac{(x+1)^2}{x} \cdot \frac{x}{(x+1)^2}} = \left(1 - \frac{2}{\text{blah}} \right)^{\text{blah}}$$

BA! $\text{blah} \rightarrow \infty$ as $x \rightarrow \infty$, $\left(1 - \frac{2}{\text{blah}} \right)^{\text{blah}} \rightarrow e^{-2}$ as $\text{blah} \rightarrow \infty$, and

$$\left(\frac{x}{x+1} \right)^2 \rightarrow 1$$

$$\frac{(x^2+1)^x}{(x+1)^{2x}} = \left(1 + \frac{-2}{\text{blah}} \right)^{\text{blah}} \rightarrow (e^{-2})^1 = e^{-2}$$

10. Does the integral $\int_1^{\infty} \frac{1}{e^x - x} dx$ converge or diverge?

(Note: 'Yes' is not considered a correct answer....)

$\frac{1}{e^x - x}$ looks "like" $\frac{1}{e^x}$ (or $\frac{1}{-x}$ or...)

$$\begin{aligned} \int_1^{\infty} \frac{1}{e^x} dx &= \int_1^{\infty} e^{-x} dx = -e^{-x} \Big|_1^{\infty} \\ &= \lim_{b \rightarrow \infty} (-e^{-x} \Big|_1^b) = \lim_{b \rightarrow \infty} (-e^{-b} - (-e^{-1})) = \lim_{b \rightarrow \infty} \frac{1}{e} - \left(\frac{1}{e^b}\right) \rightarrow 0 \\ &= \frac{1}{e} < \infty. \end{aligned}$$

$$\lim_{x \rightarrow \infty} \frac{\frac{1}{e^x}}{\frac{1}{e^x - x}} = \lim_{x \rightarrow \infty} \frac{e^x - x}{e^x} = \lim_{x \rightarrow \infty} \left(1 - \frac{x}{e^x}\right) = \lim_{x \rightarrow \infty} \frac{x}{e^x} = \frac{1}{e}$$

$$= 1 - \lim_{x \rightarrow \infty} \frac{x}{e^x} = 1 - 0 = 1 \neq 0, \infty$$

L'Hopital!

So since $\int_1^{\infty} \frac{1}{e^x} dx$ converges,

$\int_1^{\infty} \frac{1}{e^x - x} dx$ converges by limit comparison