

## Math 221 Section 5

### Practice Exam 1

Exams provide you, the student, with an opportunity to demonstrate your understanding of the techniques presented in the course. So:

**Show all work.** The steps you take to your answer are just as important, if not more important, than the answer itself. If you think it, write it!

1. (20 pts.) Find the (implicit) solution to the initial value problem

$$\frac{dy}{dt} = \frac{te^{y+t}}{y} \quad y(0) = 2 .$$

2. (15 pts.) Use Euler's method with a stepsize of  $h = 1$  to approximate the solution to the initial value problem

$$y' = ty^3 - t^3y \quad y(0) = 1$$

at time  $t = 3$  .

How would we alter this problem to find a better approximation to  $y(3)$  ?

3. (20 pts.) Find the general solution to the differential equation

$$\frac{dy}{dx} = e^{x^2} - \frac{1}{x}y$$

for  $x > 0$  .

4. (20 pts.) A vat initially contains 300 liters of a salt solution with a concentration of 4 grams/liter. Solution with a concentration of 5 grams/liter is flowing into the vat at a rate of 2 liters/minute, while the (well-stirred) contents of the vat are draining off at a rate of 3 liters/minute. What will the concentration of solution in the vat be when the vat contains 200 liters of solution?

5. (25 pts.) A new model of population growth predicts that

$$\frac{dp}{dt} = kp - (a + bp)p^2$$

(i.e., rising population will lead to ever increasing death rates due to “unnatural causes”). Observation suggests that the correct constants for the equation are  $k = 6$ ,  $a = 1$ ,  $b = 1$ . Sketch the direction field for this equation (hint: paying attention to the horizontal tangents will help), and find the (implicit) solutions to the resulting differential equation. Describe the behavior of the solution with  $p(0) = 3$ , for large  $t$ ; what limit, if any, does  $p(t)$  tend to?.

[On the original exam, a figure to sketch the field on was provided.]