

Math 314/814 Matrix Theory
Solutions to some of the Exam 2 practice problems

2. Find a basis for \mathbb{R}^3 from among the vectors

$$\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 2 \\ 1 \end{bmatrix}, \text{ which } \underline{\text{includes}} \text{ the vector } \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}.$$

In order to ensure that $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ is included, we can write it as the first column of a matrix

$$A = \begin{bmatrix} 1 & 1 & 3 & -1 \\ 1 & 2 & 2 & 2 \\ 1 & 3 & 1 & 1 \end{bmatrix}, \text{ and row reduce:}$$

$$\begin{bmatrix} 1 & 1 & 3 & -1 \\ 1 & 2 & 2 & 2 \\ 1 & 3 & 1 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 3 & -1 \\ 0 & 1 & -1 & 3 \\ 0 & 2 & -2 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 3 & -1 \\ 0 & 1 & -1 & 3 \\ 0 & 0 & 0 & -4 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 3 & -1 \\ 0 & 1 & -1 & 3 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

so we have pivots in the 1st, 2nd, and 4th columns, and so the corresponding columns of A ,

$$\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} -1 \\ 2 \\ 1 \end{bmatrix},$$

form a basis for $\text{col}(A)$. But since there are 3 of them, they also form a basis for \mathbb{R}^3 .

3. Suppose $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is a linear transformation, and suppose that

$$T \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \\ 3 \end{bmatrix}, T \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 3 \\ -2 \\ 1 \end{bmatrix}, \text{ and } T \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ -3 \\ 5 \end{bmatrix}.$$

What is $T \begin{bmatrix} 2 \\ 3 \\ -5 \end{bmatrix}$?

$$\begin{bmatrix} 2 \\ 3 \\ -5 \end{bmatrix} = 2 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 3 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + (-5) \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}, \text{ so, since } T \text{ is linear,}$$

$$T \begin{bmatrix} 2 \\ 3 \\ -5 \end{bmatrix} = T(2 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 3 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + (-5) \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}) = 2T \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 3T \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + (-5)T \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} =$$

$$2 \begin{bmatrix} -1 \\ 2 \\ 3 \end{bmatrix} + 3 \begin{bmatrix} 3 \\ -2 \\ 1 \end{bmatrix} + (-5) \begin{bmatrix} 1 \\ -3 \\ 5 \end{bmatrix} = \begin{bmatrix} -2 + 9 - 5 \\ 4 - 6 + 15 \\ 6 + 3 - 25 \end{bmatrix} = \begin{bmatrix} 2 \\ 13 \\ -16 \end{bmatrix}$$

4. Find the determinant of the matrix $B = \begin{bmatrix} 1 & 3 & 2 & 2 \\ 2 & 5 & 2 & 2 \\ 0 & 2 & 3 & 4 \\ 3 & 3 & 1 & 3 \end{bmatrix}$.

By row reduction:

$$\begin{bmatrix} 1 & 3 & 2 & 2 \\ 2 & 5 & 2 & 2 \\ 0 & 2 & 3 & 4 \\ 3 & 3 & 1 & 3 \end{bmatrix} \xrightarrow{(1)} \begin{bmatrix} 1 & 3 & 2 & 2 \\ 0 & -1 & -2 & -2 \\ 0 & 2 & 3 & 4 \\ 0 & -6 & -5 & -3 \end{bmatrix} \xrightarrow{(-1)} \begin{bmatrix} 1 & 3 & 2 & 2 \\ 0 & 1 & 2 & 2 \\ 0 & 2 & 3 & 4 \\ 0 & -6 & -5 & -3 \end{bmatrix} \xrightarrow{(1)} \begin{bmatrix} 1 & 3 & 2 & 2 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 7 & 9 \end{bmatrix} \xrightarrow{(-1)} \begin{bmatrix} 1 & 3 & 2 & 2 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 7 & 9 \end{bmatrix} \xrightarrow{(1)} \begin{bmatrix} 1 & 3 & 2 & 2 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 9 \end{bmatrix} \xrightarrow{(1)}$$

ending in a diagonal matrix, so

$$\det(B) = (1)(-1)(1)(-1)(1)[(1)(1)(1)(9)] = 9.$$

Or, by expanding along the 1st column:

$$\begin{aligned} \det(B) &= (1) \begin{vmatrix} 5 & 2 & 2 \\ 2 & 3 & 4 \\ 3 & 1 & 3 \end{vmatrix} - (2) \begin{vmatrix} 3 & 2 & 2 \\ 2 & 3 & 4 \\ 3 & 1 & 3 \end{vmatrix} + (0) \begin{vmatrix} 3 & 2 & 2 \\ 5 & 2 & 2 \\ 3 & 1 & 3 \end{vmatrix} - (3) \begin{vmatrix} 3 & 2 & 2 \\ 5 & 2 & 2 \\ 2 & 3 & 4 \end{vmatrix} \\ &= (1)((5) \begin{vmatrix} 3 & 4 \\ 1 & 3 \end{vmatrix} - (2) \begin{vmatrix} 2 & 2 \\ 1 & 3 \end{vmatrix} + (3) \begin{vmatrix} 2 & 2 \\ 3 & 4 \end{vmatrix}) - (2)((3) \begin{vmatrix} 3 & 4 \\ 1 & 3 \end{vmatrix} - (2) \begin{vmatrix} 2 & 2 \\ 1 & 3 \end{vmatrix} + (3) \begin{vmatrix} 2 & 2 \\ 3 & 4 \end{vmatrix}) - \\ &\quad (3)((3) \begin{vmatrix} 2 & 2 \\ 3 & 4 \end{vmatrix} - (5) \begin{vmatrix} 2 & 2 \\ 3 & 4 \end{vmatrix} + (2) \begin{vmatrix} 2 & 2 \\ 2 & 2 \end{vmatrix}) \\ &= (1)[(5)(5) - (2)(4) + (3)(2)] - (2)[(3)(5) - (2)(4) + (3)(2)] - (3)[(3)(2) - (5)(2) + (2)(0)] \\ &= (1)[25 - 8 + 6] - (2)[15 - 8 + 6] - (3)[6 - 10 + 0] \\ &= (1)[23] - (2)[13] - (3)[-4] \\ &= 23 - 26 + 12 \\ &= 9 \end{aligned}$$

(Other sequences of row reductions, and other expansions, are also possible.)

5. The matrix $A = \begin{bmatrix} 3 & -8 & 3 \\ 2 & -14 & 6 \\ 2 & -31 & 14 \end{bmatrix}$ has characteristic polynomial $\chi_A(\lambda) = -(\lambda + 1)(\lambda - 2)^2$.

Find the eigenvalues of A and bases for each of the corresponding eigenspaces. Is A diagonalizable? Why or why not?

Eigenvalues are solutions to $\chi_A(\lambda) = \det(A - \lambda I) = -(\lambda + 1)(\lambda - 2)^2 = 0$, so $\lambda = -1$ or $\lambda = 2$.

E_{-1} :

$$A - (-1)I = \begin{bmatrix} 4 & -8 & 3 \\ 2 & -13 & 6 \\ 2 & -31 & 15 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 & 3/4 \\ 2 & -13 & 6 \\ 2 & -31 & 15 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 & 3/4 \\ 0 & -9 & 9/2 \\ 0 & -27 & 27/2 \end{bmatrix} \rightarrow$$

$$\begin{bmatrix} 1 & -2 & 3/4 \\ 0 & 1 & -1/2 \\ 0 & -27 & 27/2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 & 3/4 \\ 0 & 1 & -1/2 \\ 0 & 0 & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & -1/4 \\ 0 & 1 & -1/2 \\ 0 & 0 & 0 \end{bmatrix}$$

so $x - z/4 = 0$, $y - z/2 = 0$, so $4x = z = 2y$, so, e.g., $\begin{bmatrix} 1 \\ 2 \\ 4 \end{bmatrix}$ is a basis for E_{-1} .

E_2 :

$$A - 2I = \begin{bmatrix} 1 & -8 & 3 \\ 2 & -16 & 6 \\ 2 & -31 & 12 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -8 & 3 \\ 0 & 0 & 0 \\ 0 & -15 & 6 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -8 & 3 \\ 0 & 0 & 0 \\ 0 & 1 & -2/5 \end{bmatrix} \rightarrow$$

$$\begin{bmatrix} 1 & 0 & -1/5 \\ 0 & 0 & 0 \\ 0 & 1 & -2/5 \end{bmatrix}$$

so $x - z/5 = 0$, $y - 2z/5 = 0$, so $5x = z$, $5y = 2z$, so e.g., $\begin{bmatrix} 1 \\ 2 \\ 5 \end{bmatrix}$ is a basis for E_2 .

Since \mathbb{R}^3 does not have a basis consisting of eigenvectors for A (or: the geometric multiplicity of $\lambda = 2$ is $1 < 2$ = the algebraic multiplicity of $\lambda = 2$), A is not diagonalizable.