

Math 325 Problem Set 1

Starred (*) problems are due Friday, August 31.

1. Let $S = \{x \in \mathbb{R} \mid x^2 + x = 0\}$ and $T = \{x \in \mathbb{R} \mid x^2 + x < 5\}$.

(*) (a) Write S and T as (small) unions of points and/or intervals.

(*) (b) Decide whether each of the following statements is true, and (briefly) explain:

$$S \subseteq \mathbb{N} ; S \subseteq T ; T \cap \mathbb{Q} \neq \emptyset ; -2.8 \in \mathbb{Q} \setminus T .$$

(c) Describe the set $U = \{x \in \mathbb{R} \mid x^2 + x < 0\}$ as a union of intervals.

2. Starting with a set S , we can construct a new set $P(S)$, the power set of S , consisting of all subsets of S . For example, $P(\{1, 2\}) = \{\emptyset, \{1\}, \{2\}, \{1, 2\}\}$.

(a) Find $P(\{1, 2, 3\})$.

(b) Show that if $S \subseteq T$, then $P(S) \subseteq P(T)$.

(c) If we set $N_k = \{1, 2, \dots, k\}$, explain why $P(N_{11})$ has twice as many elements as $P(N_{10})$.

3. Let L be the (linear) function $L(x) = ax + b$, where a and b are (real) constants and $a \neq 0$.

(*) (a) Explain why L is both one-to-one and onto.

(*) (b) Find a formula for the inverse function $M = L^{-1}$, and show that
 $L \circ M(x) = M \circ L(x) = x$ for every $x \in \mathbb{R}$.

4. Suppose the $f : A \rightarrow B$ and $g : B \rightarrow C$ are both functions.

(a) Show that if f and g are both one-to-one, then $g \circ f : A \rightarrow C$ is one-to-one.

(*) (b) Show that if $g \circ f$ is onto, then g is onto.