

# Math 445 Number Theory

September 24, 2004

Last time we found the result "*If  $p$  is an odd prime then  $x^2 \equiv -1 \pmod{p}$  has a solution  $\Leftrightarrow p \equiv 1 \pmod{4}$* " useful. Now we will explore such equations more generally.

When does the equation  $x^n \equiv a \pmod{m}$  have a solution?

We will find it useful to first deal with the warm-up problem *When does  $nx \equiv a \pmod{m}$  have a solution?* For this, we have  $nx \equiv a \pmod{m} \Leftrightarrow m|nx - a \Leftrightarrow a = nx - my$  for some  $x, y \Leftrightarrow (n, m)|a$ . Further, if  $nx_0 \equiv a \pmod{n}$ , then a complete set of incongruent solutions is given by (setting  $k = (n, m)$ )

$$x_0, x_0 + \frac{m}{k}, \dots, x_0 + (k-1)\frac{m}{k}, \quad \text{since } m|n\frac{m}{k} = m\frac{n}{k}$$

So there are in fact  $(n, m)$  solutions, if there are any.

Turning now to the main question,  $(*) x^n \equiv a \pmod{m}$ , we begin by supposing  $m$  is prime, so that there is a primitive root  $r \pmod{m}$ , i.e.,  $\text{ord}_m(r) = m-1$ . Then either  $m|a$  (so  $a \equiv 0$  and  $x = 0$  solves  $(*)$ ) or  $(a, m) = 1$ . In the latter case,  $a = r^s$  for some  $s$ . Since  $(a, m) = 1$ , any possible solution to  $(*)$  must have  $(x, m) = 1$ , as well, and so we can write  $x = r^t$  for some  $t$ . So the equation that we *really* wish to solve is

$$(**) (r^t)^n \equiv r^s \pmod{m} \quad (\text{where we wish to solve for } t).$$

But this means we wish to solve  $(r^{nt-s}) \equiv 1 \pmod{m}$ , which, since  $\text{ord}_m(r) = m-1$ , means  $m-1|nt-s$ , i.e.,  $nt \equiv s \pmod{m-1}$ . But as we have just seen, this has a solution (and we know how many)  $\Leftrightarrow (n, m-1)|s$ . Translating this back into information about  $a$ , we find that  $s = (n, m-1)q$  so  $a = r^s = r^{(n, m-1)q}$ , so, mod  $m$ ,

$$a^{\frac{m-1}{(n, m-1)}} = (r^{(n, m-1)q})^{\frac{m-1}{(n, m-1)}} = r^{(m-1)q} = (r^{m-1})^q \equiv 1^q = 1$$

Conversely, if  $a^{\frac{m-1}{(n, m-1)}} \equiv 1$ , then  $r^{s\frac{m-1}{(n, m-1)}} \equiv 1$ . Therefore  $\text{ord}_m(r) = m-1|s\frac{m-1}{(n, m-1)}$ , so  $(m-1)\frac{s}{(n, m-1)} = (m-1)y$ , so  $\frac{s}{(n, m-1)} = y$  is an integer. So  $(n, m-1)|s$ , which means  $(**)$  has a solution, and we can follow the argument back up from there to see that  $(*)$  has a solution. So we find:

If  $m$  is prime and  $(a, m) = 1$ , then

$$x^n \equiv a \pmod{m} \text{ has } \begin{cases} (n, m-1) \text{ solutions,} & \text{if } a^{\frac{m-1}{(n, m-1)}} \equiv 1 \\ 0 \text{ solutions,} & \text{if } a^{\frac{m-1}{(n, m-1)}} \not\equiv 1 \end{cases}$$

Specializing to  $n = 2$ , we have Euler's Criterion:

If  $m$  is an odd prime and  $(a, m) = 1$ , then

$$x^2 \equiv a \pmod{m} \text{ has } \begin{cases} 2 \text{ solutions,} & \text{if } a^{\frac{m-1}{2}} \equiv 1 \\ 0 \text{ solutions,} & \text{if } a^{\frac{m-1}{2}} \equiv -1 \end{cases}$$

So for example, by checking that  $13^2 = 169 \equiv -1 \pmod{17}$ , so  $13^8 \equiv 1 \pmod{17}$ , we find that  $x^2 \equiv 13 \pmod{17}$  has (two) solutions.