

# Math 445 Number Theory

October 25, 2004

**Continued fractions:** or, what happens when we “re-interpret” the Euclidean algorithm.

To compute  $(a, b)$ , we write  $a = bq_1 + r_1$ ,  $b = r_1q_2 + r_2$ , and repeat;  $r_i = r_{i+1}q_{i+2} + r_{i+2}$ , until  $r_n = 0$ . Then  $r_{n-1} = (a, b)$ . But if  $(a, b) = 1$  (so the last equation is  $r_{n-2} = 1 \cdot q_n + 0$ ) and we rewrite these calculations as

$$\frac{a}{b} = q_1 + \frac{r_1}{b}, \frac{b}{r_1} = q_2 + \frac{r_2}{r_1}, \dots, \frac{r_i}{r_{i+1}} = q_{i+2} + \frac{r_{i+2}}{r_{i+1}}, \frac{r_{n-2}}{1} = q_n + 0$$

then we can use them to express  $\frac{a}{b}$  as a *continued fraction*:

$$\frac{a}{b} = q_1 + \frac{r_1}{b} = q_1 + \frac{1}{\frac{b}{r_1}} = q_1 + \frac{1}{q_2 + \frac{r_2}{r_1}} = \dots = q_1 + \cfrac{1}{q_2 + \cfrac{1}{q_3 + \cfrac{1}{\dots + \cfrac{1}{q_n}}}}$$

For simplicity of notation, we will denote this continued fraction as  $\langle q_1, q_2, \dots, q_n \rangle$  or  $[q_1, q_2, \dots, q_n]$ , depending on whether or not we want to use the same notation as the book, or as everybody else on the planet. These continued fraction expansions are called *simple*, because the numerators are all 1, and the denominators are all positive integers. A more general theory need not require this.

And there is no reason to limit this to rational numbers! If we use the Euclidean algorithm to “compute” the gcd of  $x \in \mathbb{R}$  and 1, we would compute

$$x = 1 \cdot a_0 + r_0, \text{ i.e., } a_0 = \lfloor x \rfloor, r_0 = x - a_0 = x - \lfloor x \rfloor$$

$$1 = r_0a_1 + r_1, \text{ i.e., } \frac{1}{r_0} = a_1 + \frac{r_1}{r_0} \text{ with } a_1 \in \mathbb{N} \text{ and } r_1 < r_0, \text{ i.e., } a_1 = \lfloor \frac{1}{r_0} \rfloor, r_1 = \frac{1}{r_0} - \lfloor \frac{1}{r_0} \rfloor$$

$$\text{and, in general, } a_i = \lfloor \frac{1}{r_{i-1}} \rfloor, r_i = \frac{1}{r_{i-1}} - \lfloor \frac{1}{r_{i-1}} \rfloor$$

and we write  $x = [a_0, a_1, \dots, a_{n-1}, a_n + r_n] = [a_0, a_1, \dots, a_{n-1}, a_n + \dots]$ . For irrational numbers  $x$ , the process will not terminate. The finite continued fractions  $x_n = [a_0, a_1, \dots, a_{n-1}, a_n]$  are called the *convergents* of  $x$ .

For example, if we apply this to  $x = \sqrt{13}$ , we find

$$\begin{aligned} a_0 &= \lfloor \sqrt{13} \rfloor = 3, r_0 = \sqrt{13} - 3, & a_1 &= \lfloor \frac{1}{\sqrt{13} - 3} \rfloor = \lfloor \frac{\sqrt{13} + 3}{4} \rfloor = 1, r_1 = \frac{\sqrt{13} + 3}{4} - 1 = \frac{\sqrt{13} - 1}{4}, & a_2 &= \lfloor \frac{4}{\sqrt{13} - 1} \rfloor = \\ &\lfloor \frac{\sqrt{13} + 1}{3} \rfloor = 1, r_2 = \frac{\sqrt{13} + 1}{3} - 1 = \frac{\sqrt{13} - 2}{3}, & a_2 &= \lfloor \frac{3}{\sqrt{13} - 2} \rfloor = \lfloor \frac{\sqrt{13} + 2}{3} \rfloor = 1, r_2 = \frac{\sqrt{13} + 2}{3} - 1 = \frac{\sqrt{13} - 1}{3}, & a_3 &= \\ &\lfloor \frac{3}{\sqrt{13} - 1} \rfloor = \lfloor \frac{\sqrt{13} + 1}{4} \rfloor = 1, r_3 = \frac{\sqrt{13} + 1}{4} - 1 = \frac{\sqrt{13} - 3}{4}, & a_4 &= \lfloor \frac{4}{\sqrt{13} - 3} \rfloor = \lfloor \frac{\sqrt{13} + 3}{1} \rfloor = 6, r_4 = \frac{\sqrt{13} + 3}{1} - 6 = \frac{\sqrt{13} - 3}{1} \\ &= r_0, \end{aligned}$$

and then the process will repeat. So,  $\sqrt{13} = [3, 1, 1, 1, 1, 6, 1, 1, 1, 6, \dots] = [3, \overline{1, 1, 1, 1, 6}]$ .