

Math 872 Algebraic Topology

Problem Set # 8

Starred (*) problems due Thursday, April 13

- 39.** Giving the n -simplex $X = \Delta^n$ its standard Δ -complex structure, show that the k -skeleton of X has homology

$$\tilde{H}_i(X^{(k)}) = \begin{cases} \mathbb{Z}^{r(k,n)} & \text{if } i = k \\ 0 & \text{otherwise} \end{cases}$$

where $r(k, n) = \binom{n}{k+1}$.

(Hint: our computations preliminary to the proof of “cellular = singular” will help.)

- (*) **40.** Show that if the sequence

$$0 \rightarrow C_n \rightarrow C_{n-1} \rightarrow \cdots \rightarrow C_1 \rightarrow C_0 \rightarrow 0$$

is exact, then $\sum (-1)^i \text{rank } C_i = 0$. (Hint: pretend it's a chain complex...)

- (*) **41.** Show that if $\{\mathcal{U}, \mathcal{V}\}$ is an open cover of X and \mathcal{U}, \mathcal{V} and X all have $\oplus_i H_i$ (blah) of finite rank, then $\chi(X) = \chi(\mathcal{U}) + \chi(\mathcal{V}) - \chi(\mathcal{U} \cap \mathcal{V})$.

- 42.** Show that if \tilde{X} is an n -sheeted covering space of the finite CW-complex X , then $\chi(\tilde{X}) = n\chi(X)$. Conclude that the only non-trivial finite group that can act on an even-dimensional sphere S^{2k} without fixed points is \mathbb{Z}_2 . (Skip the hard part: the quotient by the group action is a CW-complex...)

- 43.** Show that $H_i(X \times S^n) \cong H_i(X) \oplus H_{i-n}(X)$ for every i ; here $H_i(X) = 0$ if $i < 0$.

[One approach: show that $H_i(X \times S^n) \cong H_i(X) \oplus H_i(X \times S^n, X \times D_+^n)$ (Problem #30 will help), and that $H_i(X \times S^n, X \times D_+^n) \cong H_{i-1}(X \times S^{n-1}, X \times D_+^{n-1})$ by excision and the long exact sequence of the triple $(X \times D_+^{n-1}, X \times S^{n-1}, X \times D_-^n)$.

Hatcher, p.158, # 36 gives a different approach.]