

## Math 872 Problem Set 1

Starred (\*) problems are due Thursday, January 29.

1. Show that the cone on a circle  $cS^1 = S^1 \times I / \sim$ , where  $(x, 1) \sim (y, 1)$  for  $x, y \in S^1$ , with the quotient topology, is homeomorphic to the unit disk  $D^2 = \{\vec{x} \in \mathbb{R}^2 : \|\vec{x}\| \leq 1\}$ .
2. If  $x_0 \in A \subseteq X$  and  $\pi_1(X, x_0) = \{1\}$ , show that, for any space  $Y$ , if a cts map  $f : A \rightarrow Y$  extends to a map  $g : X \rightarrow Y$ , then  $f_* : \pi_1(A, x_0) \rightarrow \pi_1(Y, f(x_0))$  is the trivial map.
- (\*) 3. [Hatcher, p.38, #5] Show that  $\pi_1(X, x_0) = \{1\}$  for every  $x_0 \in X$  if and only if every cts map  $f : S^1 \rightarrow X$  extends to a map  $F : D^2 \rightarrow X$  (i.e., for some  $F$ , we have  $F \circ \iota = f$ ).
4. [Hatcher, p.38, #2] Show that, in general, if  $\alpha, \beta : I \rightarrow X$  are paths which are *homotopic rel endpoints*,  $\alpha(0) = \beta(0) = x_0$ ,  $\alpha(1) = \beta(1) = x_1$ , then their associated change of basepoint maps are equal:  $\alpha_* = \beta_* : \pi_1(X, x_0) \rightarrow \pi_1(X, x_1)$ .

[N.B.: Hatcher calls these maps  $\beta_\alpha$  and (sorry...)  $\beta_\beta$ , for reasons I don't understand. Maybe  $\beta =$  'basepoint'?]

5. [Hatcher, p.38, #3] Show that for a path-connected space  $X$ ,  $\pi_1(X)$  is abelian  $\Leftrightarrow$  the change of basepoint maps are all independent of path, i.e.,  
for  $\alpha, \beta : I \rightarrow X$  with  $\alpha(0) = \beta(0) = x_0$  and  $\alpha(1) = \beta(1) = x_1$ , we always have  $\alpha_* = \beta_* : \pi_1(X, x_0) \rightarrow \pi_1(X, x_1)$ .

[Useful notation: our hypothesis can be expressed symbolically by saying that  $\alpha, \beta$  are maps of triples;  $\alpha, \beta : (I, 0, 1) \rightarrow (X, x_0, x_1)$ .]

- (\*) 6. Show that every homomorphism  $\varphi : \mathbb{Z} \rightarrow \pi_1(X, x_0)$  can be realized as the induced homomorphism  $\varphi = f_*$  of a continuous map  $f : (S^1, (1, 0)) \rightarrow (X, x_0)$ . [Hint: Look at  $\varphi(1) \in \pi_1(X, x_0)$ .]
- (\*) 7. [Hatcher, p.39, # 13] If  $x_0 \in A \subseteq X$  and  $A$  is path-connected, show that the inclusion-induced map  $\iota_* : \pi_1(A, x_0) \rightarrow \pi_1(X, x_0)$  is surjective  $\Leftrightarrow$  for every path  $\gamma : I \rightarrow X$  with endpoints in  $A$ ,  $\gamma(0), \gamma(1) \in A$ ,  $\gamma$  is homotopic rel endpoints to a path in  $A$ , that is,  $\gamma \simeq \alpha$  rel  $\partial I$  with  $\alpha : I \rightarrow A \subseteq X$ .
8. A *topological group* is a space  $G$  with continuous maps  $G \times G \rightarrow G$  and  $G \rightarrow G$ , denoted  $(g, h) \mapsto g \cdot h$  and  $g \mapsto g^{-1}$ , which (together with an  $e \in G$ ) make  $G$  a group. Show that for loops  $\alpha, \beta : (I, \partial I) \rightarrow (G, e)$ , the loop  $\gamma(t) = \alpha(t) \cdot \beta(t)$  is homotopic, rel endpoints, to both  $\alpha * \beta$  and  $\beta * \alpha$ . Conclude that for any topological group  $G$ ,  $\pi_1(G, e)$  is abelian.

[Hint:  $\alpha * \beta(t) = (\alpha * c_e)(t) \cdot (c_e * \beta)(t)$ .]