

### Math 872 Problem Set 3

Starred (\*) problems are due Thursday, February 19.

17. [Hatcher, p.19, # 14] Given  $v, e, f > 0$  with  $v - e + f = 2$ , build a cell structure on  $S^2$  with  $v$  0-cells,  $e$  1-cells, and  $f$  2-cells.
18. For a CW complex  $X$ , show that if the 1-skeleton  $X^{(1)}$  of  $X$  is path-connected, then  $X$  is path connected.
- (\*) 19. [Hatcher, p.20 #22] Show that if  $X$  is a CW complex,  $A, B \subseteq X$  are subcomplexes,  $A \cup B = X$ , and  $A, B$  and  $A \cap B$  are contractible, then  $X$  is contractible.
20. [Hatcher, p.20, #28] Show that if the pair  $(X, A)$  satisfies the homotopy extension property, then for any map  $f : A \rightarrow Y$  the pair  $(Z, Y)$ , where  $Z = X \amalg Y / \{a \sim f(a) : a \in A\}$  is the space built by gluing  $X$  to  $Y$  along  $A$  using  $f$ , also satisfies the homotopy extension property.
- (\*) 21. [Hatcher, p.53, #7] Let  $X$  be the quotient space formed from the 2-sphere  $S^2$  by identifying the north and south poles. Put a cell structure on  $X$  and use this to compute  $\pi_1(X)$ .
22. [Hatcher, p.54, # 14 (sort of)] Let  $X =$  the space obtained from a cube  $J^3 = J \times J \times J$ ,  $J = [-1, 1]$ , by gluing opposite square faces to one another with a 90-degree righthand twist (e.g., glue  $J \times J \times \{0\}$  to  $J \times J \times \{1\}$  by the map  $(x, y, 0) \mapsto (y, -x, 1)$ ). Describe a CW structure for  $X$  and compute a presentation for  $\pi_1(X)$ .
23. Find a cell structure for, and compute a presentation for the fundamental group of, the space  $X$  obtained by gluing two Möbius bands  $I \times I / \{(t, 0) \sim (1 - t, 1) : t \in I\}$  along their boundary circles.
- (\*) 24. [Hatcher, p.53, # 11 (sort of)] For a map  $f : X \rightarrow X$ , the *mapping torus*  $T_f$  of  $f$  is the space  $X \times I / \{(x, 0) \sim (f(x), 1) : x \in X\}$  obtained by gluing the ends of the ‘cylinder’  $X \times I$  together using  $f$ . Find a presentation of  $\pi_1(T_f)$  in terms of  $f_* : \pi_1(X, x_0) \rightarrow \pi_1(X, x_0)$  in the case when  $X = S^1 \vee S^1$  (joined along the basepoint  $x_0$ ), and  $f(x_0) = x_0$ . [Hint: treating  $T_f$  as  $X \vee S^1$  with cells attached can streamline this.]